

Archives of Pathology and Clinical Research

Volume - 3, Issue - 1

Research Article

Published Date:-2019-12-24 00:00:00

[Pregnancy and childbirth in women aged 40 and over at befelatanana maternity, Madagascar](#)

Objective: Pregnancy after age 40 remains a concern as it exposes to particular obstetrical complications. Our study aims to determine the risks of complications related to pregnancy and childbirth of women aged 40 and over.

Study design: We carried out a cross-sectional analytical study of a historical cohort comparing the progress and the outcome of pregnancy in women 40 years of age and over to those aged 20 and 35 who gave birth at the University Hospital of Obstetrics and Gynecology in Befelatanana, from 1st January 2010 to 31 December 2013.

Results: The prevalence of childbirth among 40 years old and over was 0.61%. They were multiparous and large multiparous in 88% of the cases. The analysis showed that parturients aged 40 years and older were at significant risk for caesarean section, with three times the risk of emergency caesarean section. The frequency of this emergency caesarean section increased with parity ((RR = 3.04 [2.15-4.30], p = 10-10). Among their neonates, 23.42% were hypotrophic, 22.86% premature, 12% asphyxiated at birth, 13.14% admitted to neonatal resuscitation and 5.71% died in utero, but without significant difference with the group unexposed. Perinatal death was 7.43% in women aged 40 and over vs. 4% in 20 to 35 year olds ((RR = 1.85 [0,89-3,86]; p 0.052).

Conclusion: We found that pregnancies after 40 years were not exposed to pregnancy-related pathologies or specific fetal complications. The use of an emergency cesarean is, however, frequent.

Letter to Editor

Published Date:-2019-12-03 00:00:00

[Effect evaluation of vitamin D level amongst patients with chronic hepatitis B](#)

Vitamin D has immunomodulatory and antifibrotic properties, and therefore used for treatment of many of chronic liver disease [1]. Although there are many reports on the relationship between serum 25-hydroxyvitamin D3 levels and chronic liver diseases, but the relationship between hepatitis B virus e antigen (HBeAg) and vitamin D level is still unclear.

The modification and prevention of vitamin D deficiency needs an accurate illustration of the current position in each region. Vitamin D level in patients with HBV is relatively an important issue, which has been studied in many researches. As different papers published in national and international journals.

Research Article

Published Date:-2019-07-23 00:00:00

[Longitudinal and transverse dimensions of hypertrophied cardiomyocytes in human left ventricle \(LV\) at autopsy: Heart failure might result from changes in chamber geometry without contractile weakening](#)

The shapes and sizes of human cardiomyocytes are accessible to systematic observation under most circumstances only at autopsy. This constraint has seriously curbed the study of these topics, thereby leaving a crippling gap in our understanding of heart failure. In recent years the only published ongoing findings have come from this laboratory. This article is a condensation of these reports, using those sources to develop fresh analyses designed to construct a set of organizing principles. The data are entirely retrospective thereby forbidding hypothesis testing and permitting only hypothesis formation. The hypotheses generated in this way are novel and surprising. In spite of the severe limitations in this methodology it seems possible that some useful new directions of inquiry might evolve from pursuing these original observations.

[Rosai-Dorfman disease presenting as a breast mass](#)

Rosai-Dorfman disease (RDD) is an idiopathic, benign proliferation of histiocytes that can be present in multiple organs such as lymph node, skin, soft tissue, orbit, central nerve system and bone; however, it rarely occurs in the breast. In general, RDD is a painless, firm and poorly defined lesion, which can radiologically mimic a breast cancer and is therefore an important differential consideration. The diagnosis of breast RDD is challenging, especially on a small biopsy specimen. We report a unique case of breast RDD with a literature review including common presentation, differential diagnosis, and recommended management. A high index of suspicion for this rare entity is essential to render a correct diagnosis, thereby avoiding unnecessary diagnostic tests and treatment.

[Prostatic inflammatory myofibroblastic tumor: 2-case report with literature review](#)

Inflammatory myofibroblastic tumor is rarely seen in the prostate. Most of prostatic tumors are benign. However, local aggressively invading the adjacent organs and recurrence are frequent. Rarely, the malignant transformation and metastasis occurs, especially in elderly patient. The definitive diagnosis in a small biopsy is challenging. The correct diagnosis preoperatively will help to avoid unnecessary radical prostatectomy. Interval follow up is recommended. Here we reported two cases of prostatic inflammatory myofibroblastic tumor with literature review.
